



NUCLEAR STRESS TEST - CARDIOLITE

★ **IMPORTANT - ***PLEASE REMEMBER***:**

- **NO** caffeine or decaffeinated products for 12 Hours prior to the exam.

This includes COFFEE, DECAF COFFEE, TEA, DECAF TEA, CHOCOLATE and SODA. Water is best option.

- Fast 2 HRS before the exam.

Expect to be at our facility for approximately 3 hours.

*** INSTRUCTIONS ***

- Bring a snack, drink, and any medications you may need to take with you. You will be instructed when you are able to have them.
- To walk on the treadmill please wear comfortable clothing, closed toe shoes (sneakers, etc). LADIES: **NO ONE PIECE DRESS. NO underwire bra (sports bra preferred)**
- A driver is not necessary.
- You **MUST** skip the following medication as noted below for the time frame listed by the category of medication:

BETA BLOCKERS – HOLD FOR 48 HOURS PRIOR

Cartrol (Cartelol)	Kerlone (Betaxolol)	Toprol XL (Metoprolol)	Inderal (Propranolol)
Coreg (Carvedilol)	Levatiol (Penbutolol)	Trandate (Labetolol)	Bystolic (Nebivolol)
Corzide (Nadolol)	Lopressor (Metoprolol)	Visken (Pinolol)	Blocadron (Timolol)
Inderide LA (Propranolol)	Normodyne (Labetolol)	Zebate (Bisoprolol)	Bystiloc (Timolol)
Tenoretic (Atenolol)	Ziac (Bisoprolol)	Tenormin (Atenolol)	

CALCIUM CHANNEL BLOCKERS – HOLD 36 HRS PRIOR

Cardizem (Diltiazem)	Isoptin (Verapamil)
Cardizem CD (Diltiazem)	Tiaza (Diltiazem)
Covera HS (Verapamil)	Verelan (Verapamil)
Dilacor (Diltiazem)	Tarka (Verapamil)

NITRATES – HOLD FOR 6 HRS PRIOR

Imudur (Isosorbide)	Isordil (Isosorbide)	Nitrodur Patch
---------------------	----------------------	----------------

Any questions feel free to contact us at (407) 846-0626 extension 829.



PRUEBA DE ESFUERZO NUCLEAR- (CARDIOLITE)

★ IMPORTANTE - *RECUERDE POR FAVOR***:**

- **NO** cafeína ni productos descafeinados por 12 Hours antes del examen. Esto incluye el CAFE, CAFE DESCAFEINADO, TE, TE DESCAFEINADO, CHOCOLATE Y SODA. Agua es la mejor opción.

- Ayune 2 HRS antes del examen.
- Prepárese para estar en nuestra facilidad por 3 horas que es la duración del examen.

***** INSTRUCCIONES *****

- Traiga una merienda, bebida, y cualquier medicamento que tenga que tomar en el horario que dura el examen. Le estaremos dando instrucciones a que hora las puede consumir una vez este en el stress lab.
- Para caminar en la trotadora favor de traer ropa cómoda, zapatos cerrados (tennis, etc).
DAMAS: NO UTILIZAR TRAJES DE UNA PIEZA ni brassiere con varilla.
- NO NECESITA CHOFER.
- Si consume algunos de los medicamentos mencionados abajo, **DEBERA** aguantarlos por el tiempo mencionado bajo la categoría del medicamento

BETA BLOCKERS – HOLD FOR 48 HOURS PRIOR

Cartrol (Cartelol)	Kerlone (Betaxolol)	Toprol XL (Metoprolol)	Inderal (Propranolol)
Coreg (Carvedilol)	Levatiol (Penbutolol)	Trandate (Labetolol)	Bystolic (Nebivolol)
Corzide (Nadolol)	Lopressor (Metoprolol)	Visken (Pinozolol)	Blocadron (Timolol)
Inderide LA (Propranolol)	Normodyne (Labetolol)	Zebate (Bisoprolol)	Bystilol (Timolol)
Tenoretic (Atenolol)	Ziac (Bisoprolol)	Tenormin (Atenolol)	

CALCIUM CHANNEL BLOCKERS – HOLD 36 HRS PRIOR

Cardizem (Diltiazem)	Isoptin (Verapamil)
Cardizem CD (Diltiazem)	Tiaza (Diltiazem)
Covera HS (Verapamil)	Verelan (Verapamil)
Dilacor (Diltiazem)	Tarka (Verapamil)

NITRATES – HOLD FOR 6 HRS PRIOR

Imudur (Isosorbide)	Isordil (Isosorbide)	Nitrodur Patch
---------------------	----------------------	----------------